

UPPER ENDOSCOPY (EGD)

IMPORTANT: Please read this now to be prepared for your procedure. If you have any questions, or need to cancel or postpone your procedure, please call us at **508-210-5806**.

<p>Where to report for your procedure: <u>275 Sandwich Street Plymouth, MA 02360</u></p> <ul style="list-style-type: none"> • Use Parking Lot A, which is closest to the Endoscopy Department by Urann Lobby and Valet is available.
<p>Day/ Date: _____ Arrival Time: _____</p>
<p>Procedure Time: _____ Approximate discharge time: _____</p>
<p>Doctor who will do your test: _____</p>

Advance preparation for your procedure

- ❖ **Medications** – If you take Pepto-Bismol or any iron supplement, please stop taking 7 days prior to your procedure. You can continue to take aspirin or similar pain medicines. If you take **blood-thinning medicine**, please call your prescribing physician if you did not discuss it when you booked the appointment. **Do NOT make any changes** to these medicines on your own.
- ❖ **Diabetes** – Please contact the doctor who manages your diabetes before making any changes to your diet, medications or insulin pump.
- ❖ **If you may be pregnant** – Please consult your OB/GYN doctor prior to your appointment to discuss optimal timing for your procedure, and the best sedation/anesthesia approach.
- ❖ **Sedation** – Most patients receive sedatives for procedures. Please let us know in advance about any of the following items, which could have an impact on any sedation you may receive:
 - Allergic reaction or other problems related to sedatives or pain medicine/narcotics
 - If your weight is over 300 pounds
 - You are currently taking narcotic pain medicine
 - You have severe liver disease
- ❖ As you recover from the sedatives, do not go back to work or school, make important decisions or provide care for children. You may resume all activities the next day unless otherwise instructed.
- ❖ **ARRANGE A RIDE HOME**– A responsible adult must come up to the procedure area when you are ready for discharge. You must provide us with their name and phone number when you 1st check-in. Inability to provide this information will delay or cancel your procedure. For your safety, public transportation such as taxi, bus, Uber, or Lyft is **not allowed**. No exceptions are made unless you plan to undergo the procedure without sedation. If you wish to do this, you should discuss it with your doctor in advance. **You may not drive yourself home after sedation.**

Preparing for your EGD

An upper endoscopy is an effective and safe procedure for diagnosing a variety of problems that cause symptoms in the upper part of the gastrointestinal (GI) tract. This includes the esophagus (swallowing tube), the stomach, and the duodenum (the first part of the small intestine). An upper endoscopy is sometimes called an esophagogastroduodenoscopy, or EGD.

The day before your EGD

- **At Midnight:** Stop eating all solid food. No hard candy or gum. You may drink clear liquids up until four (4) hours before your procedure.

The day of your EGD

- **4 hours before your scheduled procedure time:**
 - Stop drinking all fluids and do not take anything by mouth. (food, drink, candy, or gum). You may take any usual morning medications with a small sip of water. If you have diabetes or take blood thinners, please follow your doctor's advice regarding any changes to your medications that may be needed.

For patients taking GLP-1 agonists (Wegovy, Ozempic, Trulicity, Saxenda, Mounjaro, ect..) for weight loss, Diabetes, or any other indication please read the following:



IMPORTANT

- For patients with daily dosing hold your GLP-1 medication on the day of the procedure.
- For patients on weekly dosing hold your GLP-1 medication **1 week prior** to procedure.

IMPORTANT!

Please read at least a week before your procedure and call us if you have any questions.



Clear liquids include:

- Water
- Light-colored sodas
- Tea or coffee
- Black only- no cream or non-dairy creamers of any kind
- Clear juices
 - white grape, apple and cranberry (with no red/purple dye)
- Broth & bouillon
- Jell-O (no red)
- Popsicles (no red)

Clear liquids do **NOT** include:

- Solid food
- Gum or hard candy
- Drinks with red dye
- Dairy products
- Juice with pulp
- Alcohol